arXiv:1210.5514v1 [cond-mat.mes-hall] 19 Oct 2012

A proposal to probe quantum non-locality of Majorana fermions in tunneling experiments

Jay D. Sau[1], Brian Swingle[1], and Sumanta Tewari[2]
[1]*Department of Physics, Harvard University, Cambridge, MA 02138*
[2]*Department of Physics and Astronomy, Clemson University, Clemson, SC 29634*

Topological Majorana fermion (MF) quasiparticles have been recently shown to exist in semiconductor quantum wires with proximity induced superconductivity and a Zeeman field. Although the experimentally observed zero bias tunneling peak and a fractional ac-Josephson effect can be taken as necessary signatures of MFs, neither of them constitutes a sufficient "smoking gun" experiment. Since one pair of Majorana fermions share a single conventional fermionic degree of freedom, MFs are in a sense fractionalized excitations. Based on this fractionalization we propose a tunneling experiment that furnishes a nearly unique signature of end state MFs in semiconductor quantum wires. In particular, we show that a "teleportation"-like experiment is not enough to distinguish MFs from accidental conventional zero energy states, but our proposed tunneling experiment, in principle, can make this distinction.

PACS numbers: 03.67.Lx, 03.65.Vf, 71.10.Pm

Majorana fermions [1] are localized particle-like neutral zero energy states that occur at topological defects and boundaries in superconductors. Zero energy simply means that they occur at the chemical potential. A MF creation operator is a hermitian second quantized operator $\gamma^\dagger = \gamma$ which anti-commutes with other fermion operators. The hermiticity of MF operators implies that they can be construed as particles which are their own anti-particles [1–3]. The key issues at this time in the condensed matter context are two fold, first, we must predict and characterize materials supporting Majorana fermions and second, we must detect them experimentally. In this paper we address the second issue of experimental detection by proposing a nearly sufficent experimental signature for Majorana fermions.

MFs have recently been proposed to exist in the topologically superconducting (TS) phase of a spin-orbit (SO) coupled semiconductor 2D thin film [4, 5] or 1D nanowire [5–7] with proximity induced s-wave superconductivity and Zeeman splitting from a sufficiently large magnetic field. In principle, the MFs in such systems may be detected either by measuring the zero-bias conductance peak (ZBCP) from tunneling electrons into the end MFs [5, 8, 9] or by detecting the predicted fractional ac Josephson effect [6, 7, 10–12]. The semiconductor Majorana wire structure, which will be the system of our focus, is of particular present interest since there is experimental evidence for both the ZBCP [13–16] and the fractional ac Josephson effect in the form of doubled Shapiro steps [13].

Despite their conceptual simplicity, neither the ZBCP nor the fractional ac-Josephson effect experiments constitute a sufficient proof of MFs at the ends of topological superconducting wires. A non-quantized $(2e^2/h)$ zero bias peak, such as that observed in the recent experiments [14–16], can in principle arise even without end state MFs [17–19]. Similarly, while the observation of doubled Shapiro steps are consistent with the TS state [20–23], a fractional ac-Josephson effect can exist even in Josephson junctions made of ordinary quasi-1D p-wave superconductors such as organic superconductors [11] or the non-topological phase of the semiconductor nanowire [24]. Given these caveats as well as the considerable complexity of existing experiments, there have been several alternative proposals to detect the presence of MFs [25–28]. We emphasize that no single experiment can be expected to give a definite answer, so it is valuable to explore additional signatures that strengthen the case of MFs. Based on the inherent quantum non-locality of MFs [10, 26, 29–31], in this paper we propose an alternative tunneling experiment on semiconductor Majorana wires that furnishes a nearly sufficient signature of end-state MFs.

Non-locality arises in MFs because they differ from conventional complex (Dirac) fermions in that they have no occupation number associated with them. To define a quantum state of a system with MFs we must consider a pair of MFs. Take the pair of MFs γ_a and γ_b at the ends a and b of a nanowire (NW) shown in Fig. 1 and construct zero-energy complex fermion operator $d^\dagger = \frac{1}{2}[\gamma_a + i\gamma_b]$ associated with the pair of MFs γ_a and γ_b [10]. The quantum state of the system is then determined by the eigenvalue of $n_d = d^\dagger d = 0, 1$. Since n_d is related to the MFs by

$$n_d = \frac{1 + i\gamma_a\gamma_b}{2}, \tag{1}$$

we see that the state of the whole system is determined by non-local correlations between the fractionalized MFs γ_a and γ_b. Our central concern is how to probe this non-locality to provide a robust and sufficient criterion of MFs.

An immediate idea involves trying to inject an electron into γ_a and retrieve it from γ_b. By connecting leads to the left and the right ends of the TS wire in Fig. 1, one could imagine that an electron injected into the end a flips the occupation number n_d from $n_d = 0$ to $n_d = 1$. The injected electron can then escape from the end b flipping the state back from $n_d = 1$ to $n_d = 0$. Such a process where an electron can enter from one end a and exit at the other lead b, can be interpreted as a transfer of an electron, which we will refer to as Majorana-assisted electron tunneling. However, as has been discussed in previous works [29, 32], such a transfer occurs in a way so as to not violate locality and causality.

The amplitude for the Majorana-assisted electron tunneling [29, 30] can be written as

$$S_{a,b}^{(g,g)}(\tau) = \langle g|\gamma_a(\tau)\gamma_b(0)|g\rangle, \tag{2}$$

where $|g\rangle$ is the initial state of the Majorana wire, τ is the time-interval between the tunneling events and $\gamma_{a,b}$ are the Majorana fermion operators at the left and right end of the wire. More generally the tunneling amplitude from the end a to the end b, while the state of the nanowire system changes from g to g', can be denoted by $S_{a,b}^{(g,g')}(\tau)$. When the state $|g\rangle$ is a ground state of the TS system with a definite fermion-parity F_P, the amplitude $S_{a,b}^{(g,g)}(\tau) = -iF_P$ is non-zero. In this case the amplitude $S_{a,b}^{(g,g)}(\tau)$ can be detected by a so-called coincidence measurement for the absence of an electron on the left lead and the presence of an electron on the right lead after the tunneling event [29].

However, a coincidence measurement does not directly imply a non-zero $S_{a,b}^{(g,g)}(\tau)$ in more general situations. The amplitude $S_{a,b}^{(g,g)}(\tau)$ in Eq. 2, reflects the amplitude for being able to transfer an electron from a to b while leaving the state $|g\rangle$ invariant. On the other hand, the measurement of the coincidence probability, P_c, does not keep track of the internal state of the system. For a general system (i.e. one that may be topological or non-topological), P_c for an electron entering at a and exiting at b can be written more generally as

$$P_c = \sum_{g_1,g_2} |\langle g_2|\gamma_b\gamma_a^\dagger|g_1\rangle|^2, \tag{3}$$

where g_1, g_2 are the internal states of the wire, which are not necessarily identical. Here for a non-topological system $\gamma_{a,b}^\dagger$ are conventional fermionic operators. While TS systems with MFs have a non-degenerate ground state in a given fermion parity sector, more general systems with zero-energy Dirac end states may have multiple allowed values for g_1, g_2. Therefore, the coincidence probability P_c cannot be considered a unique signature for a topological system.

An important example of the inequivalence of P_c and $S_{a,b}^{(g,g)}(\tau)$, to which we will return later, is a non-topological superconductor with a complex fermion zero mode at each end (such a system could arise because we have an even number of weakly coupled Majorana fermions at each end). The quantum state is characterized by the occupancy n_a, n_b of the two conventional zero energy end modes. We can easily have $P_c \neq 0$ in this non-topological setup. Suppose the initial state is $g_1 \equiv (n_a = n_b = 0)$, then the sum for P_c in Eq. 3 would have a non-zero contribution from $g_2 \equiv (n_a = n_b = 1)$. The tunneling of an electron from the lead into the zero-mode at a changes the occupation from $n_a = 0$ to $n_a = 1$. On the other hand, the electron required to change the occupation of the state b from $n_b = 0$ to $n_b = 1$ comes from breaking of a Cooper pair. The other electron from the broken Cooper pair is emitted into the lead in the vicinity near b. Note that the process conserves the number of electrons within the system and cannot be eliminated even by the introduction of a finite charging energy [31]. Therefore in order to clearly distinguish this case from the process of Majorana-assisted electron tunneling (that also returns a non-zero P_c), we require $S_{a,b}^{(g,g)}(\tau)$ given in Eq. 2 itself to be non-zero. In other words, we require that the system return to the same state g after the tunneling process, so the same electron that enters at a leaves at b. In this paper we focus on how $S_{a,b}^{(g,g)}(\tau)$ can be directly measured to result in a unique signature of end state MFs [34].



FIG. 1. Schematic experiment to detect Majorana-assisted electron tunneling between the MFs a and b along the dashed lines. The electrons introduced in the system through the probe are transported via the Majorana-assisted tunneling process along the dashed line, which enclosed the flux Φ. The energy levels of these electrons, which oscillate as a function of flux Φ, are measured by the tunnel probe. By coupling the nanowires NW to the bulk superconductor SC through a Josephson junction JJ one can introduce a gate-controllable charging energy proportional to the fermion-parity.

As we now argue, the tunneling conductance experiment proposed in Fig. 1 only receives contributions from electron-tunneling processes where $S_{a,b}^{(g,g)}(\tau) \neq 0$ and therefore can be used to measure $S_{a,b}^{(g,g)}(\tau)$. The setup in Fig. 1 consists of an external quantum dot labeled

QD that is connected to the ends a and b through tunnel barriers. The gate potentials in the vicinity of QD are arranged so that a single confinement level at energy ε is present near the fermi level μ_p of the probe, which in turn is chosen to be close to the chemical potential of the electrons in the superconductor, μ_{SC}. The tunneling amplitudes $t_{a,b}$ of the tunnel barriers connecting the state in QD to a, b respectively are chosen to be smaller than the energy separation between the energy levels of the QD, so that only a single QD level participates. This is possible even for substantially long quantum dots made of InAs because of the small effective masses of InAs. In the case where the nanowire system NW in the TS phase there are MF modes at the ends a and b. In this work we will restrict ourselves to the case where the system is well-described by the non-number conserving proximity-induced superconducting Hamiltonian, so that the effects of phase fluctuations, which might limit the velocity of the Majorana assisted tunneling process, are ignored.

The physical picture is that the electron in the QD tunnels into the pair of MFs and effectively creates an interferometer such that the electron path encloses a flux Φ through the loop. Thus our set-up effectively implements a measurement of non-local correlation of MFs in nanowires, which is related to interferometric proposals to detect non-locality in MFs in topological insulators [31]. However, the proposal in Fig. 1 is more suitable for implementation in semiconductor nanowire structures that have been realized in experiments. In the case that there is a finite tunneling amplitude to go from the left end a of the TS system to the right end, the energy level of the electron in the wire can be expected to develop an $2\Phi_0 = hc/e$ periodic dependence on flux similar to the Aharonov-Bohm oscillations of the energy levels of an electron in a metallic ring. The probe, which is connected to the quantum dot QD by a weak tunneling matrix element, is used to measure the shift in the quasiparticle energy level through a peak in the tunneling conductance between the probe and the superconducting lead SC.

To demonstrate the connection between the electron-transfer amplitude across the ends a and b and the energy spectrum for electrons in Fig. 1, consider a Hamiltonian for the system labelled NW to be H_0. To ensure that the analysis here applies to both the topological and non-topological cases for NW, we will label the low-energy manifold of states that are accessed by the experiment by $|n\rangle$. The TS system is coupled to the QD through a tunneling Hamiltonian written as

$$V = \sum_\alpha (t_\alpha e^{i\varphi_\alpha} f_\alpha c^\dagger + h.c), \tag{4}$$

where f_α are the fermionic states at the end of the TS system and $c^\dagger$ is the creation operator of the active fermionic state in QD. The phases φ_α are introduced by the vector potential from the flux Φ. We will choose a gauge where $\varphi_b = 0$ and $\varphi_a = \frac{\phi}{2} = 2\pi\Phi/2\Phi_0$. In the absence of the tunneling perturbation V, the energy levels of the combined TS-QD system are written as $\rho_{n,n_0} = \nu_n + n_0\varepsilon$ where $n_0 = 0, 1$ is the occupation state of the fermionic state in the QD. Here ν_n are the unperturbed energy levels of the TS and ε is the energy of the fermionic level in QD. A subset n_g of the values taken by the TS state label n describes states of the TS system with energies near the ground state.

The action of the fermion creation operators f takes the system from a ground state n_g to a different state n. Assuming that the energy level ε in the QD is not resonant with a fermionic level in the TS, the state n will have a different energy from the state n_g i.e. $\rho_{n_0,n_g} \neq \rho_{\bar{n}_0,n}$. Integrating out these intermediate excited states then yields an effective Hamiltonian

$$\begin{aligned} h_{n_0,n_g,n'_g} &= \rho_{n_0,n_g}\delta_{n_g,n'_g} \\ &+ \sum_{n,\bar{n}_0\neq n_0} \frac{\langle n_0, n_g|V|\bar{n_0}, n\rangle\langle n_0, n'_g|V|\bar{n_0}, n\rangle^*}{E-\rho_{\bar{n}_0,n}}, \end{aligned} \tag{5}$$

in the manifold of low-energy state n_g, n'_g with energy $E \approx \rho_{n_0,n_g}$. Note that for the perturbation theory to be valid it is necessary for the gate-tuned energy ε to be tuned far off-resonance from quasiparticle excitation energies in the TS wire by more than the tunneling matrix elements that depend on t_α.

The flux-dependence of the quasiparticle energy of QD measured in the ground state is written as

$$\begin{aligned} E_{qp}(\Phi) - E_{qp}(0) &= E_{n_0=1,n_g}(\Phi) - E_{n_0=0,n_g}(\Phi) \\ &- E_{n_0=1,n_g}(0) + E_{n_0=0,n_g}(0). \end{aligned} \tag{6}$$

Using Eq. 4 the flux-dependent part of the energy for the $n_0 = 1$ QD state is determined from diagonalizing the Hamiltonian

$$\begin{aligned} \delta h_{n_g,n'_g} &= \int d\tau e^{i(E-\varepsilon)\tau}[|t_a|^2 S^{(n_g,n'_g)}_{aa}(\tau) + |t_b|^2 S^{(n_g,n'_g)}_{bb}(\tau) \\ &+ t^*_a t_b S^{(n_g,n'_g)}_{ab}(\tau)e^{-i\phi/2} + t^*_b t_a S^{(n_g,n'_g)}_{ba}(\tau)e^{i\phi/2}], \end{aligned} \tag{7}$$

where $S^{(n_g,n'_g)}_{\alpha\beta}(\tau) = \langle n_g|f_\beta(0)f^\dagger_\alpha(\tau)|n'_g\rangle$ is a generalization of the electron-tunneling amplitude defined in Eq. 2, to the case where the wire has multiple states.

In the case where the NW system is topological with the operators $f_{a,b}$ being Majorana fermions, there is a unique ground state n_g in a given fermion parity sector FP. The Hamiltonian δh in Eq. 7 is a scalar, so that the flux-dependent part of the energy can be written as

$$\delta E(\Phi) \sim \int d\tau e^{i(E-\varepsilon)\tau}[e^{i\phi/2}S^{(g,g)}_{a,b}(\tau) + e^{-i\phi/2}S^{(g,g)*}_{a,b}(-\tau)], \tag{8}$$

where $\phi = 2\pi\Phi/\Phi_0$. Thus, the presence of a finite tunneling amplitude $S^{(g,g)}_{a,b}(\tau)$ defined in Eq. 2 leads to a $2\Phi_0$ phase contribution to the energy δE.



FIG. 2. (a) Conductance as a function of probe voltage V_p and flux Φ in the presence of end MFs at a and b shows $2\Phi_0 = \frac{hc}{e}$ periodic oscillations. (b) Conductance as a function of probe voltage V_p and flux Φ in the absence of end MFs at a and b shows $\Phi_0 = \frac{hc}{2e}$ periodic oscillations.

In the case where the quantum state of the wire is associated with the fermion parity $F_{P,a}$ and $F_{P,b}$ of conventional zero-energy states at the left and right ends respectively, the quantum state is flipped (for example) from $n_g = (F_{P,a} = 1, F_{P,b} = 1)$ to $n'_g = (F_{P,b} = -1, F_{P,a} = -1)$. As a result, only the off-diagonal in fermion parity matrix element terms in Eq. 7 are non-zero. This allows one to transform the $\phi/2$ phase-dependence to a ϕ phase-dependence by a gauge transformation. The resulting spectrum in the non-topological case is only Φ_0-periodic. Thus the $2\Phi_0$ periodicity of the spectrum that may be observed for MFs is a signature of the fractionalization of the fermion parity operator F_P in a non-local way into fermionic operators defined at the ends [33].

The most direct way to measure the quasiparticle level E is by performing tunnel spectroscopy using a tunnel probe P (shown in Fig. 1) as has been done in previous experiments. In the limit of weak coupling between the probe P the conductance measures the quasi-particle energy level E and in principle the flux dependent part δE. This leads to a peak in the conductance at the quasiparticle energy that would be measured from the probe P and whose energy would depend on the applied flux with a periodicity of $2\Phi_0$ as seen in Fig. 2(b). In the absence of MFs, the superconductor can only transport electrons through Cooper pairs and the resulting level spectrum will be Φ_0-periodic.

However, the introduction of the tunnel probe P, introduces the possibility that the fermion-parity FP of the TS system can change by tunneling a fermion between the TS system and the probe. In fact, the fermion tunneling process that would change the quasiparticle number is also the process that leads to a tunneling current that allows E_{qp} to be measured in this approach. By solving Eq. 7, one finds that changing FP in the topological superconducting case shifts the phase dependence $\delta E(\Phi)$ by Φ_0, so that allowing FP to flip randomly would return the $2\Phi_0$ periodicity back to a Φ_0 periodicity and destroy the signature of the TS phase. The problem with FP flipping randomly can be fixed by introducing a charging energy E_C [31]. The charging energy makes the fermion-parity-changing process energetically unfavorable [31, 35]. The charging energy term $E_C = e^2/2C$, where C is the capacitance is introduced in the TS wire together with a Josephson coupling $E_J \gg E_C$ to an infinite superconducting lead M, so that the superconducting phase in the TS wire is essentially fixed apart from phase-slips [35, 36]. In this limit we find (see Supplementary material for derivation) that the effective Hamiltonian describing a pair of MFs is written as

$$H_{eff} \approx F_P\delta + i(\gamma_a(t_a c_a + t_a^* c_a^\dagger) + \gamma_b(t_b c_b + t_b^* c_b^\dagger)), \quad (9)$$

where $\delta \approx \frac{4}{\sqrt{\pi}} E_C \left(\frac{E_J}{E_C}\right)^{3/4} e^{-\frac{1}{2}\sqrt{E_J/E_C}}$ and $F_P = (-1)^N = i\gamma_a\gamma_b$ is the fermion parity operator. Here we are assuming that the wire is significantly longer than the decay length of the zero-energy end modes and at the same time short enough so that C is not infinite in which case δ will vanish.

In principle, it is possible to measure the $2\Phi_0$ periodicity of the quasiparticle energy indirectly without the probe P, which would change F_P. This can be done by measuring the periodicity of the total energy E_{tot} in a fixed fermion parity sector F_P. While measuring the total energy E_{tot} is difficult, it is possible to measure the current $I(\Phi) = \partial_\Phi E_{tot}(\Phi)$. Such a measurement in Fig. 1 is identical to the fractional Josephson effect [10]. As mentioned earlier, non-adiabaticity effects prevent the fractional Josephson effect from being an unambiguous signature for MFs [4]. Fixing the fermion parity by using a finite charging energy allows us to use a DC measurement, in turn avoid the non-adiabaticity effects and thus to uniquely determine the topological character of the MFs.

Summary and Conclusion: In this paper we have proposed a scheme for uniquely identifying the Majorana assisted non-local electron tunneling between two MFs at the ends of a wire in the TS phase. In principle, a non-local transfer of electrons in TS wires may be observable by a coincidence measurement in the absence of a charging energy of the TS system [29, 30]. An experimental proposal for such a coincidence measurement has been given by using a joint measurement by two point contact detectors [37]. The non-local transfer of electron in the presence of charging energy [31], which fixes the fermion parity, can also be measured by a non-local conductance or transconductance between the leads $\frac{dI_b}{dV_a}$, where a voltage applied to the left-lead a results in a current in the right lead b. Using results of Ref. 32 for symmetric $t_a = t_b = t$ we find that

$$\frac{dI_b}{dV_a} = \delta \frac{32V_a}{16\Gamma^2 + (\delta^2 - V_a^2)^2 + 8\Gamma^2(\delta^2 + V_a^2)}, \tag{10}$$

which clearly vanishes for $\delta \to 0$. Here $\Gamma \propto t^2$ is the lead-induced broadening of the MFs. However, as we have shown here that the Majorana assisted electron tunneling process using either a coincidence detection [29, 37] or by measuring the transconductance with a charging energy, while interesting, cannot be taken as a definitive signature of MF modes because even conventional near-zero energy states trapped near the spatially separated leads can also produce such non-local signature.

In contrast to the possible measurements of non-locality discussed in the previous paragraph, our proposed non-local correlation experiment in terms of the flux dependence of the tunneling spectrum provides a direct verification of the non-locality of MFs in TS wires. We emphasize that the non-locality of the end state MFs arises from the non-locality of the fermion parity, which is unique to topological systems and cannot be emulated by conventional systems[33]. In our proposal fermion parity is fixed by a superconducting single-electron transistor configuration. The scheme proposed in this paper where the tunneling amplitude is measured as a flux dependence of an energy level should be suitable for testing for MFs in the semiconductor wire system-based experiments.

Like the ZBCP and the fractional ac-Josephson signals, our interferometric signal will give a positive result if the system contains a single MF at each end. However, as we mentioned in the introduction, both the ZBCP and the ac-Josephson signal can be mimicked by non-topological physics. In our current proposal, considering the non-topological situation with two MFs at each end of a wire (as in Ref. 28), the system can in principle be prepared in a non-local initial state so that our current proposal gives a positive result. However, as shown in the supplementary section, decoherence induced by coupling to the leads disfavors the non-local initial state and washes out any signature of non-locality in the present experiment. Hence, although a weakly coupled pair of MFs (or a conventional zero-energy state) at each end may still show a ZBCP at or near zero bias at finite temperature, our interference measurement will correctly demonstrate that the system is smoothly connected to a non-topological phase.

J. D. S acknowledges support from the Harvard Quantum Optics Center. S. T. would like to thank DARPAMTO, Grant No. FA9550-10-1-0497 and NSF, Grant No. PHY-1104527 for support. B. G. S. is supported by a Simons Fellowship through Harvard University. We acknowledge valuable conversations with Bertrand Halperin and Liang Fu.

[1] E. Majorana, Nuovo Cimento **14**, 171 (1937).
[2] F. Wilczek, Nature Physics **5**, 614 (2009).
[3] M. Frantz, Physics **3**, 24 (2010).
[4] Jay D. Sau, R. M. Lutchyn, S. Tewari, S. Das Sarma, Phys. Rev. Lett. **104**, 040502 (2010).
[5] J. D. Sau, S. Tewari, R. Lutchyn, T. Stanescu and S. Das Sarma, Phys. Rev. B **82**, 214509 (2010).
[6] R. M. Lutchyn, J. D. Sau, S. Das Sarma, Phys. Rev. Lett. **105**, 077001 (2010).
[7] Y. Oreg, G. Refael, F. V. Oppen, Phys. Rev. Lett. **105**, 177002 (2010).
[8] K. Sengupta, I. Zutic, H.-J. Kwon, V. M. Yakovenko, S. Das Sarma, Phys. Rev. B **63**, 144531 (2001).
[9] K. T. Law, Patrick A. Lee, and T. K. Ng, Phys. Rev. Lett. **103**, 237001 (2009); K. Flensberg, Phys. Rev. B **82**, 180516 (2010); M. Wimmer, A.R. Akhmerov, J.P. Dahlhaus, C.W.J. Beenakker, New J. Phys. **13**, 053016 (2011).
[10] A. Y. Kitaev, Physics-Uspekhi **44**, 131 (2001).
[11] H.-J. Kwon, K. Sengupta, V. M. Yakovenko, The European Physical Journal B **37**, 349-361 (2004).
[12] L. Fu, C. L. Kane, Phys. Rev. B **79**, 161408(R) (2009).
[13] L. P. Rokhinson, X. Liu, J. K. Furdyna, arXiv: 1204.4212.
[14] A. Das, Y. Ronen, Y. Most, Y. Oreg, M. Heiblum, H. Shtrikman, arXiv:1205.7073.
[15] V. Mourik, K. Zuo, S. M. Frolov, S. R. Plissard, E. P. A. M. Bakkers and L. P. Kouwenhoven, Science DOI: 10.1126/science.1222360.
[16] M. T. Deng, C. L. Yu, G. Y. Huang, M. Larsson, P. Caroff, H. Q. Xu, arXiv:1204.4130.
[17] J. Liu, A. C. Potter, K.T. Law, P. A. Lee, arXiv:1206.1276.
[18] G. Kells, D. Meidan, P. W. Brouwer, arXiv:1207.3067.
[19] D. I. Pikulin, J. P. Dahlhaus, M. Wimmer, H. Schomerus, C. W. J. Beenakker, arXiv:1206.6687.
[20] F. Domnguez, F. Hassler, G. Platero, arXiv:1202.0642 (2012).
[21] D. M. Badiane, M. Houzet, and J. S. Meyer, Phys. Rev. Lett. **107**, 177002 (2011).
[22] D. I. Pikulin, Y. V. Nazarov, arXiv:1112.6368 (2011).
[23] P. San-Jose, E. Prada, R. Aguado, Phys. Rev. Lett. **108**, 257001 (2012).
[24] J. D. Sau, E. Berg, B. I. Halperin, arxiv 1206.4596(2012).
[25] L. Jiang, D. Pekker, J. Alicea, G. Refael, Y. Oreg, F. von Oppen, arXiv:1107.4102 (2011).
[26] C. V. Kraus, S. Diehl, P. Zoller, M. A. Baranov, arXiv:1201.3253 (2012).
[27] J. D. Sau, E. Demler, arXiv:1204.2537.
[28] B. Beri, N. R. Cooper, Phys. Rev. Lett. **109**, 156803 (2012).

[29] S. Tewari, C. Zhang, S. Das Sarma, C. Nayak, and D-H. Lee, Phys. Rev. Lett. **100**, 027001 (2008).

[30] G. W. Semenoff, P. Sodano, arXiv:0601261.

[31] L. Fu, Phys. Rev. Lett. 104, 056402 (2010).

[32] C. J. Bolech, Eugene Demler, Phys. Rev. Lett. 98, 237002 (2007).

[33] A. Turner, F. Pollman, E. Berg, Phys. Rev. B **83**, 075102 (2011).

[34] Strictly speaking, even in this case, it is possible to construct an entangled state g of the end occupations n_a and n_b so that $S_{a,b}^{(g,g)}(\tau) \neq 0$. However, as we will show later this entanglement is fragile and will be destroyed in the presence of dissipation. Note that, even though any residual gapless electronic conducting states in the quantum wire are also expected to yield a non-zero $S_{a,b}^{(g,g)}(\tau)$, in this case this quantity remains non-zero even when b is a point in the middle of the wire. In the case of Majorana assisted tunneling however, $S_{a,b}^{(g,g)}(\tau)$ is non-zero only when a and b are the two end points of the Majorana wire and in this sense an experiment that directly detects $S_{a,b}^{(g,g)}(\tau)$ with a and b as the two end points of the wire can be taken as a unique signature of the existence of MFs.

[35] B. van Heck, A.R. Akhmerov, F. Hassler, M.Burrello, C.W.J. Beenakker, New J. Phys. **14** 035019 (2012).

[36] F. Hassler, A. R. Akhmerov, C.-Y. Hou, and C. W. J. Beenakker, New J. Phys. **12**, 125002 (2010); J. D. Sau, S. Tewari, and S. Das Sarma, Phys. Rev. A **82**, 052322 (2010).

[37] Y. Cao, P. Wang, M. Gong, S-S Li, X-Q Li, arXiv:1208.3738 (2012).

Derivation of the effective Majorana Hamiltonian

To study the possibility of controlling the fermion parity variable n_d, we consider a BCS Hamiltonian containing a charging term on the TS wire that is characterized by a capacitance C. The TS wire is assumed to be coupled to an infinite superconducting lead M through a Josephson junction J. The resulting Hamiltonian for the system is written as

$$\begin{aligned} H = & \int dx dx' \psi_\sigma^\dagger(x) H_0(x\sigma, x'\sigma') \psi_{\sigma'}(x') \\ & + \Delta[\int dx e^{i\theta} \psi_\uparrow^\dagger(x) \psi_\downarrow^\dagger(x) - e^{-i\theta} \psi_\uparrow(x) \psi_\downarrow(x)] \\ & - E_J \cos\theta + E_C(n + n_F/2 - n_g)^2, \end{aligned} \tag{11}$$

where H_0 is the single-particle part of the Hamiltonian representing the semiconducting wire, Δ is the proximity-induced pairing potential, E_J is the Josephson coupling between the superconductor and the wire, $E_C(n + n_F/2 - n_g)^2$ is the charging energy resulting from a global Coulomb interaction within the superconducting-island-nanowire system. Here n is the number of Cooper pair on the island, which satisfies the canonical commutation relation $[n, \theta] = i$ and n_g is the gate charge of the TS system that is controlled by the total gate voltage on the system. $n_F = \int dx \psi^\dagger(x)\psi(x)$ is the total number of fermions on the wire.

To solve the low-energy eigenstates of this problem, we first extend the Hilbert space for the variable θ to allow the wave-function to be 4π periodic. Later we will reject all solutions in the extended Hilbert space that do not satisfy the 2π periodic boundary conditions. In the extended Hilbert space, it is convenient to apply a unitary transformation $U(\theta) = e^{i n_F \theta/2}$, which transforms $\psi^\dagger \to U(\theta)^\dagger \psi^\dagger U(\theta) = \psi^\dagger e^{i\theta/2}$. The transformation $U(\theta)$ also transforms the term $n = -i\partial_\theta \to n - n_F/2$. The transformed Hamiltonian is now written as

$$\begin{aligned} H = & \int dx dx' \psi_\sigma^\dagger(x) H_0(x\sigma, x'\sigma') \psi_{\sigma'}(x') \\ & + \Delta[\int dx \psi_\uparrow^\dagger(x) \psi_\downarrow^\dagger(x) - \psi_\uparrow(x) \psi_\downarrow(x)] \\ & - E_J \cos\theta + E_C(n - n_g)^2. \end{aligned} \tag{12}$$

The Hamiltonian H separates into an H_{BCS} written as

$$\begin{aligned} H_{BCS} = & \int dx dx' \psi_\sigma^\dagger(x) H_0(x\sigma, x'\sigma') \psi_{\sigma'}(x') \\ & + \Delta[\int dx \psi_\uparrow^\dagger(x) \psi_\downarrow^\dagger(x) - \psi_\uparrow(x) \psi_\downarrow(x)], \end{aligned} \tag{13}$$

which commutes with n, θ, together with a H_{JJ} written as

$$H_J = -E_J \cos\theta + E_C(n - n_g)^2, \tag{14}$$

where $[H_J, H_{BCS}] = 0$.

In addition to H_J and H_{BCS} it is necessary to consider the fermion tunneling terms from the leads which in the transformed basis is written as

$$H_t = \int dx [t(x) e^{i\theta/2} \psi^\dagger(x) c(x) + h.c], \tag{15}$$

where $c^\dagger(x)$ is a fermion creation operator in the lead. Specifically, we will be interested in the case where the BCS Hamiltonian has low-lying (potentially Majorana) states $f_{a,b}^\dagger$ at the ends a and b. Eliminating the low-energy gapped modes one can write

$$H_t \approx \sum_{\alpha=a,b} [e^{i\theta/2}(t_{N,\alpha} f_\alpha^\dagger + t_{A,\alpha} f_\alpha) c_\alpha + h.c]. \tag{16}$$

To return to the original Hilbert space where θ is 2π periodic, it is necessary to apply the unitary transformation $U_1(\theta) = e^{-i n_d \theta/2}$, where $n_d = \sum_\alpha f_\alpha^\dagger f_\alpha$. In the case where f_α are Majorana fermions, one must define a conventional fermion $d^\dagger = f_a + i f_b$ and define $n_d = d^\dagger d$. In either case, the applied unitary transformation replaces the $e^{i\theta/2}$ dependence of H_t by an $e^{i\theta}$ dependence. The Hamiltonian for the system now becomes

$$H \approx -E_J \cos\theta + E_C(n + n_d/2 - n_g)^2. \tag{17}$$

The spectrum of the above Hamiltonian can be gauge transformed so as to shift n by an arbitrary integer. Therefore, the spectrum of H is integer periodic. The ground state shift induced by n_d is given by $\delta \approx \frac{4}{\sqrt{\pi}} E_C \left(\frac{E_J}{E_C}\right)^{3/4} e^{-\frac{1}{2}\sqrt{E_J/E_C}}$, which is derived by solving the Mathieu equation.

Assuming that $t_{a,b} \ll \sqrt{E_J E_C}$, which is the excitation gap of the phase variable or equivalently the Josephson plasma frequency, one can project into the ground state of the phase variable. Since, $E_J/E_C \gg 1$, the phase variable θ is localized near $\theta \sim 0$ and under the projection to the ground state the Franck-Condon factor $e^{i\theta}$ can be approximated to be unity.

Dephasing of non-local correlations/entanglement for conventional fermionic zero-modes

The non-local signatures of the MFs discussed previously were consequences of robust non-local correlations that were unique to MFs. In the case of conventional low-energy (even zero-energy) modes such correlations cannot be present in a robust way. In particular, even if the system accidentally reaches a non-local correlated state, coupling to either Bosonic or fermionic dissipative bath will destroy such correlations. In the present section of the appendix, we show that coupling to a Fermionic bath leads to such desctruction of non-local correlations. We accomplish this in two sub-sections. In the first subsection, we show that when placed in contact with a fermion bath in thermal equilibrium, the system reaches a unique thermal equilibrium state. In the second subsection, we show that the system in thermal equilibrium cannot have any non-local correlations of the type that might lead to the measured interference or non-local tunneling amplitude.

Equilibration in the non-interacting case

The non-interacting fermion problem can be solved in terms of

$$\rho_{ab}(t) = i\langle\gamma_a(t)\gamma_b(t)\rangle. \tag{18}$$

The Hamiltonian in terms of Majorana operators is written as

$$H = i\sum_{a,b} h_{a,b}\gamma_a\gamma_b. \tag{19}$$

The equation of motion is written as

$$\partial_t\gamma_a = h_{a,b}\gamma_b, \tag{20}$$

with $h_{a,b}$ being anti-symmetric. The time-evolution of $\gamma_a(t)$ is written as

$$\gamma(t) = G^{(R)}(t)\gamma(0), \tag{21}$$

where $G^{(R)}(t) = e^{ht}$ for $t > 0$. Expanding $G^{(R)}(t)$ in eigenstates

$$\begin{aligned} G^{(R)}(t) &= \sum_n \psi_n\psi_n^\dagger e^{i\varepsilon_n t} \\ &= \sum_n \psi_n\psi_n^\dagger \int_{-\infty}^{\infty} \frac{e^{i\omega t}}{\omega - \varepsilon_n - i\delta} = \int_{-\infty}^{\infty} \frac{e^{i\omega t}}{\omega - ih - i\delta}. \end{aligned} \tag{22}$$

The time evolution of the density matrix is written as

$$\rho(t) = G^{(R)}(t)\rho(0)G^{(A)}(t). \tag{23}$$

Assuming that we start with a density matrix that is diagonal between system (s) and bath (b) (i.e. lead), the system density matrix

$$\rho_s(t) = G_s^{(R)}(t)\rho_s(0)G_s^{(A)}(t) + G_{sb}^{(R)}(t)\rho_b(0)G_{bs}^{(A)}(t). \tag{24}$$

Assuming the system-bath structure for the Hamiltonian and defining $g_b^{(R,A)} = (\omega \mp i\delta - H_b)^{-1}$, one can write the final system Green function as

$$G_s^{(R,A)}(\omega) = (\omega \mp i\delta - H_{sb} g_b^{(R,A)}(\omega) H_{bs})^{-1}. \tag{25}$$

The system bath interaction Green function can be written as

$$\begin{aligned} G_{sb}^{(R,A)}(\omega) &= G_s^{(R,A)}(\omega) H_{sb} g_b^{(R,A)}(\omega) \\ G_{bs}^{(R,A)}(\omega) &= g_b^{(R,A)}(\omega) H_{bs} G_s^{(R,A)}(\omega). \end{aligned} \tag{26}$$

Provided the anti-hermitean part of the fermion-self-energy

$$\Sigma_s^{(R,A)}(\omega) = Im[H_{sb} g_b^{(R,A)}(\omega) H_{bs}] \tag{27}$$

does not have a null-space (i.e. a space of zero-eigenvalues) the Green function $G_s^{(R,A)}(t)$ can be assumed to decay exponentially in time. Therefore it follows from Eq. 24 that the first term must vanish. To evaluate the second term in Eq. 24, we note that from Eq. 26, it is clear that

$$G_{sb}^{(R,A)}(t) = \int d\tau G_s(t-\tau) H_{sb} g_b(\tau). \tag{28}$$

Since $G_s(t-\tau)$ is exponentially decaying, the $\Theta(\tau)$ component of g_b is not relevant and one can approximate

$$\begin{aligned} G_{sb}^{(R,A)}(t) &\approx \int d\tau G_s(t-\tau) H_{sb} e^{-h_b\tau} \\ &= \int_0^{\infty} d\tau G_s(\tau) H_{sb} e^{-h_b(t-\tau)} \end{aligned} \tag{29}$$

up to exponential factors. Applying this identity to Eq. 24 we obtain

$$\begin{aligned} &\rho_s(t) \\ &\approx \int_0^{\infty} d\tau_1 d\tau_2 G_s^{(R)}(\tau_1) H_{sb} e^{-h_b(t-\tau_1)} \rho_b(0) e^{h_b(t-\tau_2)} H_{bs} G_s^{(A)}(\tau_2) \\ &= \int_{-\infty}^{\infty} d\tau_1 d\tau_2 G_s^{(R)}(\tau_1) H_{sb} \rho_b(0) e^{h_b(\tau_1-\tau_2)} H_{bs} G_s^{(A)}(\tau_2), \end{aligned} \tag{30}$$

which is a t-independent asymptotic value. Transforming to frequency space,

$$\rho_s(t) \approx \int_{-\infty}^{\infty} d\omega f(\omega) G_s^{(R)}(\omega) H_{sb} A_b(\omega) H_{bs} G_s^{(A)}(\omega). \tag{31}$$

Considering the integrand at ω away from a pole of $g_s(\omega)$,

$$\begin{aligned} & G_s^{(R)}(\omega) H_{sb} A_b(\omega) H_{bs} G_s^{(A)}(\omega) \\ & = i G_s^{(R)}(\omega) H_{sb} \{ g_b^{(R)}(\omega) - g_b^{(A)}(\omega) \} H_{bs} G_s^{(A)}(\omega) \\ & = i G_s^{(R)}(\omega) \{ \Sigma_s^{(R)}(\omega) - \Sigma_s^{(A)}(\omega) \} G_s^{(A)}(\omega) \\ & = i \{ G_s^{(R)}(\omega) - G_s^{(A)}(\omega) \}, \end{aligned} \tag{32}$$

which can be checked using Dyson's equation. Therefore, in the absence of a protected null-space in the dissipative part of the self-energy $\Sigma_s(\omega)$, the density matrix equilibrates to the grand-canonical thermal equilibrium value

$$\rho_s(t) \approx i \int_{-\infty}^{\infty} d\omega f(\omega) \{ G_s^{(R)}(\omega) - G_s^{(A)}(\omega) \}. \tag{33}$$

Correlations in the grand-canonical thermal state

First we prove a general lemma: Consider the grand-canonical thermal partition function of a system of Majorana fermions (or fermions) which can be partitioned into two parts L and R, so that the Hamiltonian has a symmetry $\gamma_{L,a} \to -\gamma_{L,a}$ for all MFs $\gamma_{L,a}$ on the left half of the system L. Then all correlation functions $\langle \gamma_{L,a} \gamma_{R,b} \rangle = 0$. To prove this simply apply the transformation $\gamma_{L,a} \to -U \gamma_{L,a} U^\dagger$ to

$$\begin{aligned} \langle \gamma_{L,a} \gamma_{R,b} \rangle & \propto Tr[\gamma_{L,a} \gamma_{R,b} e^{-\beta H}] = Tr[U \gamma_{L,a} U^\dagger \gamma_{R,b} U e^{-\beta H} U^\dagger] \\ & = -Tr[\gamma_{L,a} \gamma_{R,b} e^{-\beta H}] = 0. \end{aligned} \tag{34}$$

Therefore all such non-local fermionic correlations must vanish in the thermal state.

As discussed before, in the topological state, the charging energy gives rise to a fermion parity dependent term in the Hamiltonian

$$H_{nl} \propto i \gamma_a \gamma_b, \tag{35}$$

which violates the conditions for the above lemma and leaves a non-zero value for this non-local correlator. Therefore the presence of the Majorana fermion term is crucial for the appearance of a non-local correlation function that can contribute to the interference. Such an interference will not happen even if the conventional end zero mode is made of a pair of Majorana fermions. This is because, as discussed in the previous sub-section, coupling Majorana fermions to leads causes them to thermalize into the grand-canonical ensemble. Once they have thermalized in this ensemble, H_{nl} violates the conditions of the lemma and generates a non-local fermion correlation. The long-range Coulomb interaction for conventional fermionic modes does not lead to the violation of the conditions of the lemma and does not lead to any long range correlation.